

# Reassessing Weight Loss Surgery For Type 2 Diabetes

Weight loss surgery is not a cure for type 2 diabetes, but it can improve blood sugar control, according to a new study published in the *British Journal of Surgery*. Whereas some previous studies have claimed that up to 80 percent of diabetes patients have been cured following gastric bypass surgery, researchers at Imperial College London found that only 41 percent of patients achieve remission using more stringent criteria.

The research was funded by the National Institute for Health Research (NIHR) Biomedical Research Centre and awarded to Imperial College Healthcare NHS Trust and Imperial College London.

Obesity is a major risk factor for type 2 diabetes. Worldwide, 80 percent of people with type 2 diabetes are overweight or obese at the time of diagnosis. Diabetes is normally treated by using insulin injections and drugs to control blood sugar. However, many diabetic patients who had stomach surgery to lose weight found that their diabetes improved, even before they had lost any weight.

Recently the American Diabetes Association pulled together a group of experts to agree on standards by which to assess whether a patient has achieved remission of diabetes. They defined complete remission as returning to normal measures of glucose metabolism without taking diabetes medication at least one year after surgery. The new study revisited previous data on 209 patients with type 2 diabetes to evaluate the effectiveness of three types of weight loss surgery using the new criteria. They found that the remission rate was 41 percent for gastric bypass, the most effective type of surgery.

"Using the new criteria, we don't get such eye-catching figures as some that have been quoted in recent years," said Dr Carel le Roux, from the Department of Medicine at Imperial College London, who led the study. "But it's clear that weight loss surgery, particularly gastric bypass, has a significant beneficial effect on glucose control. Diabetes is a chronic, multi-system disease. Stomach surgery may not mean that patients can stop taking diabetes medication, but surgery and medication together achieve better results than either treatment on its own."

**Source URL (retrieved on 01/27/2015 - 5:27am):**

<http://www.surgicalproductsmag.com/news/2012/01/reassessing-weight-loss-surgery-type-2-diabetes>