Regular Aspirin Use Could Lead To Increased Macular Degeneration

Regular aspirin use appears to be associated with an increased risk of neovascular age-related macular degeneration (AMD), which is a leading cause of blindness in older people, and it appears to be independent of a history of cardiovascular disease and smoking, according to a report published by a *JAMA Internal Medicine*, JAMA Network publication. Aspirin is one of the most widely used medications in the world and is commonly used in the prevention of cardiovascular disease, such as myocardial infarction and ischemic stroke. While a recent study suggested that regular aspirin use was associated with AMD, particularly the more visually devastating neovascular (wet) form, other studies have reported inconsistent findings. Smoking is also a preventable risk factor for AMD, the authors write in the study background.

Gerald Liew, Ph.D., of the University of Sydney, Australia, and colleagues examined whether regular aspirin use (defined as once or more per week in the past year) was associated with a higher risk of developing AMD by conducting a prospective analysis of data from an Australian study that included four examinations during a 15-year period. Of 2,389 participants, 257 individuals (10.8 percent) were regular aspirin users. After the 15-year follow-up, 63 individuals (24.5 percent) developed incident neovascular AMD, according to the results.

"The cumulative incidence of neovascular AMD among non-regular aspirin users was 0.8 percent at five years, 1.6 percent at 10 years, and 3.7 percent at 15 years; among regular aspirin users, the cumulative incidence was 1.9 percent at five years, seven percent at 10 years and 9.3 percent at 15 years, respectively," the authors note. "Regular aspirin use was significantly associated with an increased incidence of neovascular AMD."

The authors note that any decision concerning whether to stop aspirin therapy is "complex and needs to be individualized. Currently, there is insufficient evidence to recommend changing clinical practice, except perhaps in patients with strong risk factors for neovascular AMD (e.g., existing late AMD in the fellow eye) in whom it may be appropriate to raise the potentially small risk of incident neovascular AMD with long-term aspirin therapy," the authors conclude.

Source URL (retrieved on *11/26/2014 - 4:32am*):
http://www.surgicalproductsmag.com/news/2013/01/regular-aspirin-use-could-lead-increased-macular-degeneration?qt-recent_videos=0&qt-most_popular=0